



## Content

**AISHA Elfateh**

**EFFECTS OF TEN WEEKS OF INSTABILITY RESISTANCE TRAINING (BOSU BALL) ON MUSCULAR BALANCE AND THE LEARNING LEVEL OF FENCING BASICS / p. 273**

**AKSOY Cemil, SARITAŞ Nazmi, COŞKUN Betül**

**\*THE COMPARISON OF ANXIETY LEVEL BETWEEN MALE AND FEMALE ATHLETES IN AMATEUR TEAMS / p. 280**

**AYNUR Yilmaz, GONUL Tekkursun Demir, OGUZ Kaan Esenturk**

**\*HEALTH-PROMOTING LIFESTYLE BEHAVIORS OF EMPLOYEES IN PUBLIC SECTOR / p. 285**

**BÁTOROVÁ Michaela, MOTYČKA Jaroslav, ŠŤASTNÝ Jan<sup>1</sup>, LEPKOVÁ Hana**

**\*BIOMECHANICAL ANALYSIS OF FRONT CRAWL ARM STROKE AND LEG KICK BY TACHOGRAPH MEASURING SYSTEM / p. 293**

**BIANCALANA Vincenzo, NICOLA Mingozzi, ANNUNZIATA Passalia**

**\*LIFETIME SEDENTARY IS A MAJOR CAUSE OF CHRONIC DISEASES / p. 301**

**BONDOC-IONESCU Dragos, OANCEA Bogdan Marian**

**\*STUDY ON THE STREAMLINE OF THE U15 WOMENS BASKETBALL FREE THROWS / p. 306**

**BONDOC-IONESCU Dragos, NEAMTU Mircea, BONDOC-IONESCU Alexandru, MIHAI Ilie, OPREA Doru**

**\*TESTING KINEMATIC AND ERGOMETER PARAMETERS FOR PERFORMING THE PHASE CONTACT - BEAT - IMPULSE TO THE HIGH JUMP / p. 315**

**BUŢU Ioana Maria, CĂTUNĂ George Cristian, TEODORESCU Anemari Simona, IORGA Anca, GHEORGHE Georgeta Ioana**

**\*TESTING METHOD USING SPACE ORIENTATION ABILITY TO UNTRAINED PERSON MATORIN / p. 323**

**CATANESCU Andreea, CURIŢIANU Ioana**

**\*STUDY REGARDING THE DEVELOPMENT OF GENERAL AND SPECIFIC MOTOR SKILLS IN ALPINE SKIING CATEGORY U10 / p. 329**

**ÇELİK O. Burçak, KURTİPEK Serkan, İLHAN E. Levent**

**\*INCREASING BULLYING TENDENCY IN SCHOOLS: A RESEARCH IN THE CONTEXT OF EXTRACURRICULAR SPORTIVE ACTIVITIES / p. 335**

**CICMA Ioan Teodor**

**\*THE INCREASE OF THE DEVELOPMENT INDICES OF THE REACTION SPEED, USING SPECIFIC MEANS OF THE HANDBALL GAME (VII<sup>th</sup> GRADE) / p. 343**

**CICMA Ioan Teodor, MEREUŢĂ Claudiu**

**\*THE MANAGEMENT OPTIMIZATION OF THE TECHNICAL PREPARATION OF INTERMEDIATE PLAYERS, IN THE HANDBALL GAME, ECHELON JUNIORS I (GIRL'S) / p. 349**



**CIOCIOI Alexandru-Florin, MACOVEI Sabina, ZAHIU Mihaela**

**\*JOINT MOBILITY VALUES IN PRIMARY SCHOOL PUPILS BEFORE AND AFTER THE SUMMER HOLIDAY / p. 355**

**CIOROIU Silviu Gabriel, NECHITA Florentina**

**\*SWIMMING NEUROMUSCULAR CONTROL / p. 364**

**COJOCARU Adin Marian, COJOCARU Marilena**

**\*FUNCTIONAL TRAINING IN MAINTAINING THE PHYSICAL PREPARATION VOLLEYBALL PLAYER / p. 370**

**COSMA Germina, RUSU Ligia, BURILEANU Alin, NANU Marian, STĂNCESCU Camelia**

**\*THE INFLUENCE OF PHYSICAL EXERCISE ON THE CORRECTION OF DEFICIENT ATTITUDES – A CASE STUDY / p. 377**

**COSMA Germina, DRAGOMIR Marian, DUMITRU Roxana, LICĂ Eliana, GHEȚU Roberta**

**\*THE DANCE IMPACT ON THE MOTOR ABILITY IN CHILDREN / p. 382**

**DOCU AXELERAD Any, DOCU AXELERAD Daniel, FRECUȘ Corina, MUSAT George**

**\*COMPARATIVE EVALUATION OF REHABILITATION AND ACTOVEGIN TREATMENT VERSUS REHABILITATION TREATMENT IN PAINFUL DIABETIC NON INSULIN-DEPENDENT NEUROPATHY / p. 387**

**DUMITRU Mariana, MOROIANU Miruna**

**\*TEACHING STRATEGIES FOR LEARNING THE BASKETBALL GAME IN THE PRIMARY CYCLE / p. 397**

**ENE-VOICULESCU Carmen, ENE-VOICULESCU Virgil**

**\*INDICATIVE MODEL OF SYSTEM IMPLEMENTATION IN TECHNICAL PROCEDURES OF THE MAIN ALGORITHM IN SWIMMING 50M OBSTACLES IN CONSONANCE WITH THE THEME SYLLABUS – NATO / p. 398**

**FIEROIU Emil**

**\*STRATEGIES TO IMPROVE KINETIC MEANS FOR REHABILITATION REGARDING KNEE TRAUMATIC PATHOLOGY IN ATHLETES / p. 406**

**FIEROIU Emil**

**\*THE IMPORTANCE OF KINESIOTHERAPY IN THE TREATMENT AND PROPHYLAXIS OF HYPERTENSION IN ADULTS / p. 412**

**GHEORGHE Daniel, IVAN Paula**

**\*STUDY ON THE HIGH PERFORMANCES ATHLETES AT 800 m DISCIPLINE / p. 418**

**GHEORGHE Daniel, IVAN Paula**

**\*COMPARATIVE STUDY OF SOMATIC AND MOTOR CHARACTERISTICS OF THE POLE VAULT ATHLETES PARTICIPANTS AT THE LAST EIGHT EDITIONS OF THE WORLD CHAMPIONSHIPS OUTDOOR / p. 425**

**GONUL Tekkursun Demir, AYNUR Yılmaz, OGUZ Kaan Esenturk, EKREM Levent Ilhan**

**\*THE PERCEPTION OF THE HEADMASTER TOWARDS PHYSICALEDUCATION COURSE / p. 430**

**GÖNÜL Tekkursun Demir, AYNUR Yılmaz, OĞUZ Kaan Esenturk, EKREM Levent Ilhan**

**WHY DOES MY CHILD SWIM? MECHANISM OF PARENT GUIDANCE / p. 442**

**HESHAM Elgushy**



**IRON STATUS FOR THE EGYPTIAN FEMALE PLAYERS IN RUNNING COMPETITION (SHORT-MIDDLE-LONG DISTANCES) – COMPARATIVE STUDY / p. 449**

**HESHAM Elgushy**

**\*THE IMPACT OF CONCURRENT TRAINING ON CERTAIN PULMONARY, PHYSICAL VARIABLES AND RECORD LEVEL OF MIDDLE DISTANCES FOR YOUNG ATHLETICS / p. 454**

**HUSSEIN Abaza, MOHAMED Saad**

**PLAY POSITIONS AND LEFT VENTRICULAR MASS AND IT RELATIONSHIP WITH PHYSICAL VARIABLES FOR SOCCER PLAYERS / p. 462**

**IONESCU Oana-Cristiana, CORDUN Mariana**

**\*CONSIDERATIONS REGARDING THE IMPROVEMENT OF COORDINATION CAPACITY IN BLIND CHILDREN / p. 468**

**IORGA Anca, BUȚU Ioana Maria, TEODORESCU Anemari Simona, GHEORGHE Georgeta Ioana, CĂȚUNĂ George Cristian**

**\*COMPARATIVE STUDY BETWEEN ATHLETES IN RHYTHMIC GYMNASTICS AND DANCE ANATOMIC AND PHYSIOLOGIC AND DEVELOPMENT SPECIFIC FORCE / p. 477**

**ISAKOVIĆ Marko, DAMIAN Roxana**

**\*\*THE ROLE OF PREVENTIVE PRACTICE IN REDUCING THE NUMBER OF INJURIES OF HANDBALL PLAYERS / p. 484**

**ISPAS Gheorghe Andrei-Marius, MACOVEI Sabina**

**\*BENEFITS OF USING PILATES APPARATUS IN MOTOR ACTIVITIES FOR KEEPING FIT / p. 490**

**IZZO Ricardo, SOPRANZETTI Seila**

**\*WARM UP / p. 497**

**JIANU Anca**

**\*INFLUENCE OF RESPIRATORY GYMNASTICS IN ELDERLY HIPERTENSIVE SUBJECTS / p. 502**

**KENIOUA Mouloud, BACHIR Boughera, BACHA Foudil Samir**

**\*JOB SATISFACTION FOR PHYSICAL EDUCATION TEACHERS AND ITS RELATIONSHIP TO JOB PERFORMANCE AND ORGANIZATIONAL COMMITMENT / p. 507**

**KUNDERA Václav, HERBERGER Daniel**

**\*THE COOPERATION OF YOUTH AND TOP CLUBS OF SHOOTING IN THE CZECH REPUBLIC AS A BASIS FOR IMPROVING THE CURRENT STATE OF THE MEMBER BASE / p. 512**

**KURTIPEK Serkan, ÇELİK O. Burçak, YENEL İ. Fatih, İLHAN E. Levent**

**\*INVESTIGATION OF TIME MANAGEMENT SKILLS ACCORDING TO PERSONALITY CHARACTERISTICS OF UNIVERSITY STUDENTS WHO STUDY SPORT EDUCATION / p. 520**

**LOK Neslihan, LOK Sefa**

**\*CORRELATION BETWEEN DEPRESSIVE SYMPTOMS AND PHYSICAL ACTIVITY LEVELS OF ELDERS, LIVING IN A CERTAIN AREA / p. 526**

**LOK Sefa, LOK Neslihan, TASGIN Erdal**

**\*PHYSICAL ACTIVITY PREVALENCE OF HIGH SCHOOL STUDENTS / p. 532**



**MAN Maria Cristina, GANERA Cătălin**

**\*A STUDY ON ATHLETES' HEART RATE CHANGING WHILE PERFORMING A 21 DAYS TRAINING COURSE AT AN ALTITUDE OF 2000M / p. 537**

**MARINESCU Gheorghe, TICALĂ Laurențiu Daniel, DULCEAȚĂ Victor, BIDIUGAN Simona Nicoleta**

**\*EFFORT ANALYSIS IN REAL TIME DURING A FOOTBALL GAME – JUNIOR II USING GPSports DEVICE / p. 548**

**MATTEUCCI Ivana**

**\*THE PROMOTION OF WELLBEING IN ADOLESCENT MIDDLE SCHOOL STUDENTS: HIGH INTENSITY COMMUNICATION AND SELF-ESTEEM IN SCHOOL ORGANIZED SPORTS / p. 555**

**MEHTAP Bekir, TASGIN Erdal, LOK Sefa**

**\*\*COENZYME Q<sub>10</sub> AND CARDIOVASCULAR SYSTEM / p. 564**

**MEZEI Mariana, BOTA Aura**

**\*CONCEPTUAL AND PRACTICAL ASPECTS IN THE PREPARATION OF A CHOREOGRAPHIC ROUTINE IN HIGH PERFORMANCE AEROBIC GYMNASTICS – EXAMPLE OF BEST PRACTICE / p. 570**

**MUSTAFA Can Koç, NAZMI Saritaş, ŞULE Kirbaş, HÜSNIYE Bülbül**

**\*EXAMINING NUTRITIONAL HABITS OF SOCCER PLAYERS AT YOUTH DEVELOPMENT TEAMS / p. 576**

**NEGREA Valentin, MUŞAT George**

**\*DEVELOPMENT OF MOTOR SKILLS THROUGH MOVEMENT GAMES AND CONTESTS TO FIFTH GRADERS / p. 585**

**NOVAKOVA Svatava<sup>1</sup>, BLAHUTKOVA Marie<sup>2</sup>, LEPKOVA Hana**

**\*MANAGEMENT OF IMPLEMENTATION OF WATER TOURISM FOR YOUTH IN THE CZECH REPUBLIC / p. 590**

**OGUZ Kaan Esenturk, GONUL Tekkursun Demir, AYNUR Yilmaz, EKREM Levent Ilhan**

**\*EVALUATION OF THE EXTRACURRICULAR SPORTIVE ACTIVITIES FROM THE VIEW OF PHYSICAL EDUCATION TEACHERS / p. 599**

**OGUZ Kaan Esenturk, MEHIBE Akandere, ERKAN Yarimkaya, AYNUR Yilmaz**

**\*THE EFFECT OF PHYSICAL ACTIVITY PROGRAM ON TRAIT ANXIETY LEVEL ON ADULTS / p. 609**

**ONEA Gheorghe Adrian, BALINT Lorand**

**\*PRACTICAL MOTOR EXAMPLES USED BY COACHES TO OPTIMIZE THE LEARNING PROCESS FOR YOUNG HURDLES RUNNERS - A REVIEW / p. 619**

**PETREANU Manuela, BUȚU Ioana Maria, PETREANU Adrian Gheorghe**

**\*COMPARATIVE STUDY ON THE POTENTIAL BIOMOTRIC STUDENTS FROM THE UNIVERSITY "SPIRU HARET" AND U.M.F. "CAROL DAVILA" / p. 624**

**POPA Cristian, DAMIAN Mirela, POPA Corina**

**\*EDUCATIONAL ASPECTS OF HYGIENE TAUGHT IN THE PHYSICAL EDUCATION LESSON/ p. 629**

**POPESCU Daniela Corina, MIHĂILĂ Ion**

**\*THE EVALUATION PARAMETERS TRAINING SPECIFIC TO PRACTICE OF PERFORMANCE HANDBALL BY OF THE "SCALA" IT APPLICATION / p. 634**



**POPESCU Florentina, PORFIREANU Maria-Cristiana, RISTEA Cristian**

**\*THE INTEREST OF YOUTH IN SPORT ACTIVITIES / p. 641**

**RABOLU ELENA**

**\*OPERATIONAL STRATEGIES FOR SAFEGUARDING HEALTH USING KINETIC MEANS IN THE INSTITUTIONALIZED ELDERLY / p. 647**

**RADESCU Petre, TEODORESCU Silvia, URZEALĂ Constanța**

**\*STUDY ON THE NEUROMUSCULAR CONTROL ASSESSMENT BY USING THE CARTESIAN MOTION ANALYSIS / p. 652**

**RAȚĂ Bogdan Constantin, RAȚĂ MARINELA, CRĂCIUN Sergiu**

**\*PERFORMANCE PATTERNS FOR OUTDOOR EVENTS IN JUNIOR III FOR GIRLS PENTATHLON-HEXATHLON AND BOYS HEPTATHLON-OCTATHLON / p. 659**

**SABAU Elena, NICULESCU Georgeta, GEVAT Cecilia**

**\*THE PITTSBURGH SLEEP QUALITY INDEX - A MEAN OF SLEEP ASSESSING / p. 668**

**ȘTEFĂNESCU Cătălin Aurelian**

**\*DIGITAL ANALYSIS OF PLAYER'S POSITIONING AND MOVEMENT DURING A BASKETBALL OFFICIAL GAME / p. 674**

**SYED Kamaruzaman Syed Ali, MOHD Faisal Mohamed, MUHAMAD Azhar Zailani, WAIL Muil Alhaj Said Ismail**

**\*PLANNING AND PREPARING PHYSICAL EDUCATION TEACHERS ON CARDIOVASCULAR ENDURANCE, MUSCULAR ENDURANCE, AND MUSCLE STRENGTH IN THE IMPLEMENTATION OF PHYSICAL EDUCATION CURRICULUM OF FORM 4 (FITNESS STEM) / p. 680**

**TAREK Sadek**

**EFFECTS OF CUPPING THERAPY BASED ON STABILIZATION CORE EXERCISES ON LOW BACK PAIN FOR SOCCER PLAYERS IN STATE OF UNITED ARAB EMIRATES / p. 684**

**TASGIN Erdal, MEHTAP Bekir, LOK Sefa**

**\*\*EXERCISE AND HYPOPHYSIS HORMONES / p. 691**

**TASGİN Erdal, LOK Sefa**

**\*EFFECTS OF EXHAUSTIVE EXERCISE INVESTIGATION OF RATS IBUPROFEN OF ADVERSE CARDIAC EVENTS / p. 696**

**TASKIN Mine, AKKOYUNLU Yağmur**

**EFFECT OF ANAEROBIC POWER ON QUICKNESS IN WOMEN NATIONAL TAEKWONDO ATHLETES / p. 701**

**TEODORESCU Silvia, NICOARĂ Alexandru Adrian**

**\*STUDY REGARDING THE NEED TO OBJECTIFY EVALUATION IN LATIN-AMERICAN DANCES / p. 706**

**TEUȘDEA Claudiu Cristian, PETRESCU Toma**

**\*TECHNICAL DETAILS OF THE FOREHAND MEN'S TENNIS / p. 711**

**ALPHABETICAL AUTHOR INDEX / p. 721**

**Technical Requirements to Elaborate Scientific Papers / p. 723**

\* the abstract was published in the 16th I.S.C. "Perspectives in Physical Education and Sport" - Ovidius University of Constanta, May 20-21, 2016, Romania

\*\* the abstract was published in the 15<sup>th</sup> I.S.C. "Perspectives in Physical Education and Sport" - Ovidius University of Constanta, May 22-23, 2015, Romania



## ❖ ALPHABETICAL AUTHOR INDEX

### A

AISHA E. / p. 273  
AKKOYUNLU Y. / p. 701  
AKSOY C. / p. 280  
ANNUNZIATA P. / p. 301  
AYNUR Y. / p. 285, 430, 442, 599, 609

### B

BACHA F.S. / p. 507  
BACHIR B. / p. 507  
BALINT L. / p. 619  
BÁTOROVÁ M. / p. 293  
BIANCALANA V. / p. 301  
BIDIUGAN S.N. / p. 548  
BLAHUTKOVA M. / p. 590  
BONDOC IONESCUCU A. / p. 315  
BONDOC IONESCUCU D. / p. 306, 315  
BOTA A. / p. 570  
BURILEANU A. / p. 377  
BUȚU I.M. / p. 323, 477, 624

### C

CĂTĂNESCU A. / p. 329  
CĂTUNĂ G.C. / p. 323, 477  
ÇELİK O.B. / p. 335, 520  
CICMA I.T. / p. 343, 349  
CIOCIOI A.F. / p. 355  
CIOROIU S.G. / p. 364  
COJOCARU A.M. / p. 370  
COJOCARU M. / p. 370  
CORDUN M. / p. 468  
COSMA G. / p. 377, 382  
COȘKUN B. / p. 280  
CRĂCIUN S. / p. 659  
CURIȚIANU I. / p. 329

### D

DAMIAN M. / p. 629  
DAMIAN R. / p. 484  
DOCU AXELERAD A. / p. 387  
DOCU AXELERAD D. / p. 387  
DRAGOMIR M. / p. 382  
DULCEAȚĂ V. / p. 548  
DUMITRU M. / p. 392  
DUMITRU R. / p. 382

### E

EKREM L.I. / p. 430, 442, 599  
ENE-VOICULESCU C. / p. 398

ENE-VOICULESCU V. / p. 398  
ERKAN Y. / p. 609

### F

FIEROIU E. / p. 406, 412  
FRECUȘ C. / p. 387

### G

GANERA C. / p. 537  
GEVAT C. / p. 668  
GHEORGHE D. / p. 418, 425  
GHEORGHE G.I. / p. 323, 477  
GHEȚU R. / p. 382  
GONUL T.D. / p. 285, 430, 442, 599

### H

HERBERGER D. / p. 512  
HESHAM E. / p. 449, 454  
HUSSEIN A. / p. 462  
HÜSNIYE B. / p. 576

### I

İLHAN E.L. / p. 335, 520  
IONESCUCU O.C. / p. 468  
IORGA A. / p. 323, 477  
ISAKOVIĆ M. / p. 484  
ISPAS G.A.M. / p. 490  
IVAN P. / p. 418, 425  
IZZO R. / p. 497

### J

JIANU A. / p. 502

### K

KENIOUA M. / p. 507  
KUNDERA V. / p. 512  
KURTİPEK S. / p. 335, 520

### L

LEPKOVÁ H. / p. 293, 590  
LICĂ E. / p. 382  
LOK N. / p. 526, 532  
LOK S. / p. 526, 532, 564, 691, 696

### M

MACOVEI S. / p. 355, 490  
MAN M.C. / p. 537  
MARINESCUCU G. / p. 548  
MATTEUCCI I. / p. 555



MEHIBE A. / p. 609  
MEHTAP B. / p. 564, 691  
MEREUȚĂ C. / p. 349  
MEZEI M. / p. 570  
MIHAI I. / p. 315  
MIHAILĂ I. / p. 634  
MOHAMED S. / p. 462  
MOHD F.M. / p. 680  
MOROIANU M. / p. 392  
MOTYČKA J. / p. 293  
MUHAMAD A.Z. / p. 680  
MUSTAFA C.K. / p. 576  
MUȘAT G. / p. 387, 585

#### N

NANU M. / p. 377  
NAZMI S. / p. 576  
NEAMTU M. / p. 315  
NECHITA F. / p. 364  
NEGREA V. / p. 585  
NICOARĂ A.A. / p. 706  
NICOLA M. / p. 301  
NICULESCU G. / p. 668  
NOVAKOVA S. / p. 590

#### O

OANCEA B.M. / p. 306  
OGUZ K.E. / p. 285, 430, 442, 599, 609  
ONEA G.A. / p. 619  
OPREA D. / p. 315

#### P

PETREANU A.G. / p. 624  
PETREANU M. / p. 624  
PETRESCU T. / p. 711  
POPA Co. / p. 629  
POPA Cr. / p. 629  
POPESCU D.C. / p. 634  
POPESCU F. / p. 641  
PORFIREANU M.C. / p. 641

#### R

RABOLU E. / p. 647  
RADESCU P. / p. 652  
RAȚĂ B.C. / p. 659  
RAȚĂ M. / p. 659  
RISTEA C. / p. 641  
RUSU L. / p. 377

#### S

SABAU E. / p. 668  
SARITAȘ N. / p. 280  
SOPRANZETTI S. / p. 497  
ŠŤASTNÝ J. / p. 293

STĂNCESCU C. / p. 377  
SYED K.S.A. / p. 680

#### Ș

ȘTEFĂNESCU C.A. / p. 674  
ȘULE K. / p. 576

#### T

TAREK S. / p. 684  
TASGIN E. / p. 532, 564, 691, 696  
TASKIN M. / p. 701  
TEODORESCU A.S. / p. 323, 477  
TEODORESCU S. / p. 652, 706  
TEUȘDEA C.C. / p. 711  
TICALĂ L.D. / p. 548

#### U

URZEALĂ C. / p. 652

#### W

WAIL M.A.S.I. / p. 680

#### Y

YENEL İ.F. / p. 520

#### Z

ZAHIU M. / p. 355





# Technical Requirements to Elaborate Scientific Papers

Page set up:

-A4 format

-margins: top,bottom, right 1,5cm, left 2,5cm;

-footnote: authors affiliations and country, corresponding author email address;

-abstracts: 400 words maximum, preferably structured on: aim/objectives/problem statement, methods, results, conclusions, key words (3-5 words);

-full text: introduction, methods, results, discussions, conclusions, acknowledgments (optionally) references.

Full paper must be 5-10 pages, 10 font TNR, 2 columns, one space

## TITLE OF THE PAPER

SURNAME AND NAME of the paper's author or authors<sup>1</sup>

Example:

FISHER Diana<sup>1</sup>, VIOLETTE Anna<sup>2</sup>, LE BOUC Ians<sup>3</sup>

Footnotes will consist of:

a) name of department, name of institution (if necessary), name of university, city and address of university, native country (Home, Times New Roman, Size 8, Justify),

b) for the author who deals with the correspondence for the paper or reprint: name of department, name of institution (if necessary), name of university, city and address of university, native country followed by the phrase CORRESPONDENCE AND REPRINT REQUESTS: (Home, Times New Roman, Size 8, Justify, Caps Lock) name of the author, address, e-mail, phone and/or fax number (if necessary) (Home, Times New Roman, Size 8, Justify) and

c) in a new paragraph, the source of the material support in the shape of GRANT-s (if necessary)(Home, Times New Roman, Size 8, Justify), written after the phrase GRANT SUPPORT: (Home, Times New Roman, Size 8, Justify, Caps Lock).

(follow as example the footnotes)

Technical requirements to elaborate the structured informative abstract:

### Abstract

*Aim.* The aim of this study is to examine the relationship between skinfolds method (accu-measure caliper) and near-infrared method (FUTREX 1000 Personal Body Fat Tester)

*Methods.* We used Romanian university students (27 males and 97 females). The body fat percentage was measured by two methods: the skinfolds measurements...

*Results.* Body fat estimated with accu-measure caliper was moderate correlated with body fat estimated with FUTREX for women ( $r = 0.41$ )...

*Conclusions.* We cannot consider that one method of body composition analysis (skinfolds method or near-infrared method) is more accurate than...

*Keywords:* skinfolds method, near-infrared method, percentage of body fat, fat mass, free fat mass, Romanian students.

Technical requirements to elaborate the non-structured indicative or informative abstract:

<sup>1</sup> Department of Obstetrics, Gynecology and Women's Health, University of Missouri, Columbia, MO 65212, USA.

<sup>2</sup> Department of Obstetrics, Gynecology and Women's Health, Division of Biological Sciences, University of Missouri, Columbia, MO 65212, USA. CORRESPONDENCE AND REPRINT REQUESTS: Alissa Viollet, NW509 Health Sciences Center, 1 Hospital Dr., Columbia, MO 65212, USA. [aviollet@missouri.edu](mailto:aviollet@missouri.edu), tel. 573-882-6334, fax. 573-882-6399

<sup>3</sup> Department of Anatomy, Institute of Biomedical Sciences, University of São Paulo, CEP São Paulo 05508-900, Brazil.  
GRANT SUPPORT: Eunice Kennedy Shriver National Institute of Child Health and Human Development HD055231.





## Abstract

The aim of this study was to examine the relationship between skinfolds method (accu-measure caliper) and near-infrared method (FUTREX 1000 Personal Body Fat Tester) for body fat percent, fat mass and free fat mass estimations, in Romanian university students. We used Romanian university students (27 males...

**Keywords:** skinfolds method, near-infrared method, percentage of body fat, fat mass, free fat mass, Romanian students.

The page layout of the research paper is 2.5 cm from the top, bottom, left and right margins (Page Layout, Margins, Top 2.5cm, Bottom 2.5cm, Left 2.5cm, Right 2.5cm), portrait oriented (Page Layout, Orientation, Portrait), A4 (Page Layout, Size, A4 - 21cm x 29.7cm).

The title of the paper, the name of the author (authors) and the abstract will be written on a single column, following the rules of laying out the page.

The chapters: Introduction, Methods, Results, Discussions, Conclusions, Thanks (if necessary) and Bibliography will be written on two columns, except for the tables and charts which will be written on a single column, following the rules of laying out the page.

The space between the title of the paper, the name of the author or authors of the paper, abstract, introduction, methods, results, discussions, conclusions and bibliography is one line (Enter, Font Size 10); the space between the writing and the tables or charts is also one line (Enter, Font Size 10).

The titles of the sub-chapters will be written in bold (Home, Times New Roman, Size 10, Bold, Justify, First Line Indent 0,5cm). All the paragraphs will present a 0,5 cm size compared to the margin (First Line Indent 0,5cm). The text will have the following technical characteristics: Home, Times New Roman, Size 10, Justify. Between the titles of the sub-chapters and the text there will be no space.

Example of laying out the page and arranging the text:

## Relationship between skinfolds and near-infrared (FUTREX 1000) methods for body fat estimation in Romanian university students

<sup>a</sup>  
IONESCU Tudor Madalin<sup>1</sup>, MARCU Andrei<sup>2</sup>

### Abstract

**Aim.** The aim of this study was to examine the relationship between skinfolds method (accu-measure caliper) and near-infrared method (FUTREX 1000 Personal Body Fat Tester) for body fat percent, fat mass and free fat mass estimations, in Romanian university students.

**Methods.** We used Romanian university students (27 males and 97 females). The body fat percentage was measured by two methods: the skinfolds measurements (accu-measure caliper) and near-infrared measurement (Futrex 1000).

**Results.** Body fat estimated with accu-measure caliper was moderate correlated with body fat estimated with FUTREX for women ( $r = 0.41$ ) and for men ( $r = 0.55$ ). Fat mass (skinfolds method) skinfolds method and free fat mass (skinfolds method) were moderate correlated with fat mass (near-infrared method), respectively free fat mass (near-infrared method) for women ( $r = 0.41$ , respectively  $r = 0.41$ ) and correlated for men ( $r = 0.60$ , respectively  $r = 0.60$ ).

**Conclusions.** We cannot consider that one method of body composition analysis (skinfolds method or near-infrared method) is more accurate than the other because we don't apply a gold standard method of measurement, for subjects. However, near-infrared method tends to have higher estimations of body fat, then skinfolds method on Romanian students.

**Keywords:** skinfolds method, near-infrared method, percentage of body fat, fat mass, free fat mass, Romanian students.

### Introduction

The increase in obesity is a global phenomenon that is even being addressed by the World Health Organization (World Health Organization, 2003), as

well as by medical and government organizations in the world.

One of factors that contribute to body composition changes, respectively to body fat percent



grow up is physical inactivity or sedentary lives (National Institutes Of Health, 1998).

Factors, such as age, gender, level of adiposity, physical activity and ethnicity influence the choice of method and equation. To date, race-specific SKF (American Indian women, Black men, and Asian adults), BIA (American Indian women and Asian adults), and NIR (American Indian women and White women) equations have been developed (Heyward, 1996).

Infrared is not an indicator of body composition in the pre-adolescent population on an individual basis. This method continues to be no accurate, cost-effective means to assess individual body composition by a rapid, noninvasive methodology (Michael, Jan, Wendy, 2003).

Larger prediction errors have been reported with the lower cost, hand-held Futrex 1000 model. Because of these errors, the manufacturer's equations for the Futrex 1000 are not recommended to assess body composition (Wagner and Heyward, 1999).

Kamimura et al. 2009, cannot consider that one method of body composition analysis (SKF method, bioelectrical impedance analysis, or NIR method) is more accurate than the other because they didn't apply a gold standard method, for patients on long-term hemodialysis therapy. However, the most simple, long-established, and inexpensive method of SKF thickness seems to be still very useful for assessing body fat (Kamimura, Jose Dos Santos, Avesani, Fernandes Canziani, Draibe, Cuppari, 2003).

In a healthy group of 29 subjects examined by Elia et al., NIR method had little or no advantage over other simple methods in predicting body composition measured by classical whole-body densitometry. NIR method was also found to underestimate body fat increasingly as the degree of adiposity increased. This under-estimation was found to be particularly marked in a small and separate group of grossly obese women, BMI greater than 50 kg/m<sup>2</sup>, whose body composition was assessed by total body potassium as well as by densitometry (Dumitru, 1997).

Heyward et al. concluded that all three field methods, respectively SKF, bioelectric impedance and NIR compared with hydrostatic weighting, accurately estimate the percent of body fat for nonobese women; however, none of these three methods is suitable for estimating the percent of body fat for obese women (Heyward, Cook, Hicks, Jenkins, Quatrochi, Wilson, 1992).

One study concluded that, SKF is higher correlated with under water weighting than did

FUTREX 5000 with under water weighting for males (0.95 versus 0.80), females (0.88 versus 0.63), and the whole group (0.94 versus 0.81) and FUTREX 5000 overestimated body fat in lean subjects with less than 8% fat and underestimated it in subjects with greater than 30% fat. Analyzing this, the authors concluded that, SKF give more information and more accurately predict body fat, especially at the extremes of the body fat continuum (McLean and Skinner, 1992).

The present findings indicate that, the FUTREX 5000 provide more accurate estimates of body fat percent than the FUTREX 5000A or FUTREX 1000 instruments (Smith, Johnson, Stout, Housh, Housh, Evetovich, 1997). Continued research with expanded populations is needed to further demonstrate and evaluate the utility of FUTREX 5000A device (Cassady, Nielsen, Janz, Wu, Cook, Hansen, 1993).

Conway et al. concluded that, body composition (percentage fat) estimated in 53 adults (23 to 65 years of age) by infrared interactance, is correlated with SKF ( $r = 0.90$ ) measurements. They concluded that, the method is safe, noninvasive, rapid, easy to use, and may prove useful to predict percentage body fat, especially in the obese (Conway, Norris, Bodwell, 1984).

SKF method is still a reliable technique of BF estimation, but if it's not realized with the most accurately instruments the results trends to have errors in BF estimation and FM, respectively FFM (Cyrino, Okano, Glaner et al., 2003). The NIR method is still a questionable technique for BF estimation (McLean and Skinner, 1992; Michael, Jan, Wendy, 2003; Wagner and Heyward, 1999).

The objective of this study is to examine the relationship between skinfolds (SKF) method (accu-measure caliper) and near-infrared (NIR) method (FUTREX 1000 Personal Body Fat Tester) for body fat percent (BF), fat mass (FM) and free fat mass (FFM) estimation, in Romanian university students.

## Methods

The subjects were white Caucasian and students at faculties of Ovidius University in Constanta. The aims and methods of the study were explained to the participants, who chose freely to participate in this study. As a result, the sample included 127 students (97 females and 27 males), with age between 18 and 23 years old.

Body height was evaluated with an error of 0.1 centimeters and body weight was evaluated with a calibrated digital scale, with an error of 0.25 kilograms. For this measurement the subjects were dressed summarily. BMI was calculated to estimate



the category of weight for each subject by using the Quetelet formula (Dumitru, 1997).

Percent of body fat was estimated with two methods. The first method consisted in calculation of body fat percent with Jackson and Pollock, (1978), equation, for male subjects and Jackson, Pollock and Ward, (1980), equation, for female subjects. The abdominal (taken vertically with a broad grip, 5cm. lateral to the omphalion (centre of the umbilicus)), chest (taken obliquely along the natural cleavage line

of the pectoral between the axilla and nipple) and thigh (vertical fold taken midway between the inguinal crease and proximal border of the patella) skinfolds were measured for ...

.....

### Results

In table 1 the differences between sexes were significant only for body height ( $t = 9.838$ ) and body weight ( $t = 5.841$ ).

**Table 1.** Physical characteristics of the subjects

Variables	M ± SD	
	Males (n = 27)	Females (n = 97)
Age (years <sup>months</sup> )	19 <sup>7</sup> ± 0 <sup>11</sup>	20 <sup>1</sup> ± 2 <sup>8</sup>
Body height (cm)	1.789 ± 0.078 *	1.63 ± 0.059
Body weight (kg)	66.074 ± 11.135 *	52.722 ± 7.842
BMI (kg/m <sup>2</sup> )	20.598 ± 2.929	19.811 ± 2.485

\* differences between sexes,  $p < 0.05$ .  
BMI, body mass index; M, mean; SD, standard deviation; n, number of subjects.

In table 2 the differences between sexes were significant for all variables (BFskf,  $t = 13.278$ ; FMskf,  $t = 6.346$ ; FFMskf,  $t = 11.498$ ; BFnir,  $t = 7.856$ ; FMnir,  $t = 2.883$ ; FFMnir,  $t = 9.861$ ). All variables from SKF method had significant correlations with their correspondent variable from NIR method, when body height, body weight and age

were controlled. BFskf was moderate correlated with BFnir for women ( $r = 0.41$ ) and for men ( $r = 0.55$ ). FMskf and FFMskf were moderate correlated with FMnir, respectively FFMnir for women ( $r = 0.41$ , respectively  $r = 0.41$ ) and correlated for men ( $r = 0.60$ , respectively  $r = 0.60$ ).

**Table 2.** Differences between SKF method and NIR method

Variables	Skinfold method (Accu-measure caliper) M ± SD	
	Males (n = 27)	Females (n = 97)
BFskf (%)	8.962 ± 4.407 *†	21.886 ± 4.704 *
FMskf (kg)	6.25 ± 4.006 *†	11.806 ± 4.085 *
FFMskf (kg)	59.824 ± 8.207 *†	40.915 ± 4.512 *
Variables	Infrared method (Futrex 1000) M ± SD	
	Males (n = 27)	Females (n = 97)
BFnir (%)	13.074 ± 5.988 †	22.805 ± 4.475
FMnir (kg)	8.97 ± 5.431 †	12.164 ± 3.615
FFMnir (kg)	57.104 ± 8.225 †	40.557 ± 5.486

\* correlated with BFnir, FMnir and FFMnir for males, respectively for women, when height, weight and age are controlled,  $p < 0.05$ ;  
† differences between sexes,  $p < 0.05$ .  
BFskf, body fat - skinfolds method; FMskf, fat mass - skinfolds method; FFMskf, free fat mass - skinfolds method; BFnir, body fat - infrared



method; FMnir, fat mass - infrared method; FFMnir, free fat mass - infrared method; M, mean; SD, standard deviation; n, number of subjects.

.....  
**Discussion**

Compared with the anthropometric reference data 1988 – 1994 from United States (National Health and Nutrition Examination Survey, 2005), body height for our subjects was slightly higher for men and slightly lower for women, compared with the corresponding values for Americans. The body weight was lower, for both men and women, compared with the corresponding values for Americans.

.....

**Conclusions**

The conclusions must be reported directly to the hypotheses of the paper and derive directly from the chapter Discussions. The conclusions that are not fully backed-up by the data found or that are based on unjustified affirmations must be avoided. New hypotheses can be concluded or attach some recommendations, if the case be.

**Acknowledgments**

I thank all students for participating in this study. No funding was used for this study.

**References**

Cassady SL, Nielsen DH, Janz KF, Wu YT, Cook JS, Hansen JR, 1993, Validity of near infrared body composition analysis in children and adolescents, Med Sci Sports Exerc, 1993 Oct; 25(10):1185-1191.

Conway JM, Norris KH, Bodwell CE, 1984, A new approach for the estimation of body composition: infrared interactance. Am J Clin Nutr, 1984 Dec; 40(6):1123-1130.

Cyrino ES, Okano HA, Glaner FM et al., 2003, Impact of the use of different skinfold calipers for the analysis of the body composition. Rev Bras Med Esporte, 2003; 9(3):150-153.

National Institutes of Health (NIH). Clinical Guidelines On The Identification, Evaluation, And Treatment Of Overweight And Obesity In Adults. The Evidence Report. Publication No. 98-4083, 1998 Sep: XI-XXX.

.....

**Attention!!!** First of all, the article is written on a single column until it is finalized. After finalizing it, you select the whole text after the abstract until the first table or chart and you turn it into two columns. The same operation is done, in order, for (the whole) texts between charts and/or tables; also, the (whole) text, from the last table or chart until the bibliography inclusive, will be turn into two columns. **The paper must be 5-10pages.**

**Tables**

The tables including data will be done on a single column and they cannot be introduced into the text as photographs. The counting (consecutive) and the title of the table (conclusive and concise) will be written on the top right hand. The reference to the table (the quotation in the text) will be found in the text that precedes the table. The number of the table, the title of the table, the results, the statistical section and the abbreviation section will be a constitutive part of the table. It is recommended that you merge the data in as few tables as possible. The additional black lines in the tables including data will be colored in white (Table Tools, Design, Pen Color, White, urmat de Draw Table prin care se trasează peste liniile negre suplimentare culoarea albă).

Table 1. Physical characteristics of feminine subjects

Variables	Subjects with dominant upper and lower right limb(n = 8)		Subjects with dominant upper and lower left limb (n = 8)	
Height (cm.)	163,25 ± 4,95	3,032%	162,5 ± 4,309	2,652%
Weight (kg.)	66,088 ± 7,343	11,111%	67,038 ± 5,352	7,984%
IMC (kg/m <sup>2</sup> )	24,745 ± 1,827	7,383%	25,368 ± 1,439	5,673%
Percentage of body fat(%)	26,625 ± 2,873	10,791%	26,55 ± 2,964	11,164%
Fat mass (kg.)	17,739 ± 3,56	20,069%	17,91 ± 3,235	18,063%

The values are presented as M ± DS și CV%.

IMC, index of body mass; M, mean; DS, standard deviation; CV, variability coefficient; n, number of subjects.



The connection between the data in the table and the statistical section will be done through identification letters counted in alphabetical order or identification symbols used in the order \*, †, ‡, §, ||, ¶, \*\*, ††, ‡‡, etc.; inside the table, the letters or the identification symbols will be written in the superscript(Home, Superscript) immediately after the data, and inside the statistical section, the identification letters will be written before the hyphen and the statistical comments and the identification symbols immediately before the statistical comments (without a hyphen). The tables from other publications should be used with the author's (authors') permission, indicating the bibliographic source where it was taken from.

Example:  $0,851 \pm 0,044^a$

Example: a – significantly different compared to the force ratio F150 Right side flexion/ F150 Left side flexion, 0°, for the subjects who practise football, respectively athletics (triple jump),  $F(2, 12) = 5,5$ ;

Table 2. Means of results of maximum isometric force ratios for feminine subjects who practise different sports

Force ratio	Handball (n = 5)	Football (n = 5)	Athletics (triple jump) (n = 5)
F130 Flexion/ F110 Extension (30°)	$0,589 \pm 0,109$ 18,506%	$0,556 \pm 0,075$ 13,489%	$0,565 \pm 0,05$ 8,85%
F150 Right side flexion/ F150 Left side flexion (0°)	$0,851 \pm 0,044^{ab}$ 5,17%	$0,942 \pm 0,056^c$ 5,945%	$0,919 \pm 0,03^d$ 3,264%
F120 Right side rotation/ F120 Left side rotation (-30°)	$0,972 \pm 0,07$ 7,202%	$0,825 \pm 0,227$ 27,515%	$1,052 \pm 0,019^e$ 1,806%

a – significantly different compared to the mean of the force ratio F150 Right side flexion/ F150 Left side flexion, 0°, for subjects who practise football, respectively, athletics (triple jump),  $F(2, 12) = 5,5$ ;

b – significantly different compared to the mean of the force ratio F150 Right side flexion/ F150 Perfectly ballanced left side flexion (when all the force ratios are equal to 1), 0°,  $t=7,572$ ;

c – significantly different compared to the mean of the force ratio F150 Right side flexion/ F150 Perfectly ballanced left side flexion (when all the force ratios are equal to 1), 0°,  $t=2,316$ ;

d – significantly different compared to the mean of the force ratio F150 Right side flexion/ F150 Perfectly ballanced left side flexion (when all the force ratios are equal to 1), 0°,  $t=6,037$ ;

e – significantly different compared to the mean of the force ratio F120 Right side rotation/ F120 Perfectly ballanced lesft side rotation (when all the force ratios are equal to 1), -30°,  $t=6,12$ ;

The values are presented as  $M \pm DS$  and CV%; Significance limit established at  $p < 0,05$ .

M, mean; DS, standard deviation; CV, variability coefficient; n, number of subjects; t, test t student ; F, test ANOVA.

### Figures

The tables which contain figures will be done on a single column. The counting (consecutive) and the title of the figure (conclusive and concise) will be written on the bottom left side immediately after the figure. The reference to the figure (the quotation in the text) will be found in the text that precedes the table which contains the figure. The figure, the number of the figure, the title of the figure, the statistical section (if necessary) and the abbreviation section will be a constitutive part of the table that contains the figure. When symbols, numbers or letters are used to identify the parts of the figure, each of them should be explained clearly in the statistical section. It is recommended that you merge the data in as few figures as possible. The lines of the table that contains the figure will be transparent. (Table Tools, Design, Borders, No Borders).






Figure 27. The evolution of means of maximum isometric force and the degree of perception at different tests. Nm, Newton\*meter; IT, initial testing; ImedT, intermediary testing; FT, final testing.

The figures will have a resolution of minimum 250 dpi for a better understanding after the print. The figures will be presented in original sizes in the text (sizes chosen by the author(s) of the paper), not to be subsequently modified. The electronic formats accepted are: Bitmap (.bmp), JPEG (.jpg, .jpeg) or GIF (.gif). The results and the statistical explanations will be presented in one way – data in the table, figure in the table or text; these ways of presenting can be combined but they do not have to repeat themselves.

### Measures

Length, height, weight and volume will be specified in metrical units (meter, kilogram or litre or their decimal multiples). Temperature will be specified in degrees Celsius (°C). Blood pressure will be specified in mm column of mercury (mmHg). Other clinical measurements will be specified in the International System of Units (International System of Units (SI)).

### Abbreviations and symbols

The standard abbreviations must be used. You should avoid introducing abbreviations into the title or in the abstract. An abbreviation in parentheses will be preceded by the full description, only the first time the abbreviation is used in the text and only if the abbreviation is not a standard measure unit.

**Example:** Body weight, body composition, resting metabolic rate (RMR), respiratory quotient (RQ), temperature, fasting serum glucose, insulin, free fatty acids, and ghrelin were assessed at baseline and after 21 d (12-h fast) and 22 d (36-h fast) of alternate-day fasting.

RMR and RQ did not change significantly from baseline to day 21, but RQ decreased on day 22 ( $P < 0.001$ ), which resulted in an average daily increase in fat oxidation of  $\geq 15$  g.

### Bibliography

Wuthiekanun V, Chierakul W, Langa S, et al. 2006, Development of antibodies to Burkholderia pseudomallei during childhood in melioidosis-endemic northeast Thailand. Am J Trop Med Hyg 2006 Jan 12;74(10):1074-5.